

# The Miscarriage Association



---

*acknowledging pregnancy loss*

## Investigations following recurrent miscarriage

If you would like general information or to talk to someone else who has experienced recurrent miscarriage and can offer understanding and support, please contact:

The Miscarriage Association  
c/o Clayton Hospital  
Northgate  
Wakefield  
West Yorkshire WF1 3JS

Telephone: 01924 200799  
Fax: 01924 298834  
[www.miscarriageassociation.org.uk](http://www.miscarriageassociation.org.uk)

© The Miscarriage Association 2009  
Registered Charity Number 1076829 (England & Wales) SC039790 (Scotland)  
A company limited by guarantee, number 3779123  
Registered in England and Wales

RM/8/09

## Introduction

One in one hundred couples trying for a baby experiences recurrent miscarriage, usually defined as three or more miscarriages in a row. This leaflet aims to outline some of the investigations which may be carried out at that time. Although this leaflet is written for both partners, the term “you” will generally be used to mean the woman who has miscarried.

## Why investigate only after three miscarriages?

Many couples who have been through a miscarriage are anxious to find out why it happened, especially if that information can improve their chances of success in their next pregnancy. Most, however, will not be offered investigations after a single miscarriage, or even two. Although this can be distressing, this is because most women who have one or two miscarriages will go on to have a healthy pregnancy next time. This suggests that their miscarriages were due to chance rather than to an underlying cause.

If a couple has had three or more consecutive miscarriages, statistics show that there is more likely to be an underlying cause or causes and so tests are usually offered at this point. This does not necessarily mean that a cause or causes will be found. Although we are learning more about the causes of miscarriage, there is much that is still unknown.

If you do undergo investigations for recurrent miscarriage, it is possible, therefore, that many of the tests performed will be reported as normal and a clear cause for the miscarriages will not be found. This can be frustrating but it does mean that there is a very high chance of the next pregnancy being successful. If a problem **is** identified, there is still a good chance of having a successful pregnancy.

## What investigations might be offered?

This list includes tests which may be offered (we explain the medical terms on the next pages):

- Chromosomal analysis: (a) of blood from both parents (karyotype)  
(b) of the baby (fetal karyotype)
- Lupus anticoagulant and anticardiolipin antibodies (to exclude a condition known as Antiphospholipid Syndrome)
- Other immunological investigations
- Hormonal blood tests
- Weight and height
- Investigations of the anatomy of the uterus (womb) and cervix
- Tests for infection
- Research investigations

# Chromosomal analysis or karyotype: parents

## What are chromosomes?

Chromosomes carry the genetic information for each individual. Everyone has 23 pairs of chromosomes, making 46 in all. All but one pair are identical in men and women. The 23rd pair – the sex chromosomes – decide the individual's gender and are therefore different. Men normally have one X and one Y chromosome and women have two X chromosomes. A baby inherits half of its chromosomes from its mother and half from its father.

## How can chromosomes cause a problem?

About half of all miscarriages occur as a result of a chromosomal abnormality in the baby or fetus. In most cases where this abnormality causes miscarriage, the problem is not passed on from a parent, but happens when the egg and sperm meet, or early in the development of the fertilised egg. This is more common with advancing maternal age.

Between three and five percent of couples with recurrent miscarriage have a problem with a chromosomal abnormality called a **balanced translocation**. In this situation, part of the information from one chromosome is replaced by that of another chromosome. Although this doesn't cause a problem to the affected parent, it can be passed on to the baby and cause an **unbalanced** translocation, where some genetic information is present twice and some is missing. This can lead to miscarriage.

*The Miscarriage Association can provide a leaflet on balanced translocation.*

## Testing

Chromosomal analysis involves taking a blood test from both the man and the woman and sending the samples to a genetics laboratory. The results may take up to six weeks to obtain, as the cells have to be specially processed before they can be examined under the microscope.

## Treatment

There is no treatment which can alter the chromosomes in an individual if they are already abnormal. If the analysis shows that you or your partner carry an abnormality, then you will be offered specialist genetic counselling to give you more information and help you decide about future pregnancies. This may include discussion of pre-implantation genetic diagnosis (PGD); or the use of donor eggs or sperm.

# Chromosomal analysis of the baby: fetal karyotype

Your clinic may offer to carry out chromosomal analysis of fetal tissue, although this can depend on the laboratory facilities available. It involves sending tissue from the miscarriage to the genetics laboratory where it undergoes the same process as for blood. Unfortunately a result is obtained only in approximately half of cases.

It can take up to eight weeks to obtain the results. If the result is abnormal, but both parents have a normal chromosome pattern, then the abnormality in the baby is unlikely to recur in a subsequent pregnancy.

# Tests for lupus anticoagulant and anticardiolipin antibodies (Antiphospholipid or Hughes Syndrome)

## What are these?

An antibody is part of the body's defence mechanism. Lupus anticoagulant (LA) and anticardiolipin antibodies (aCL) are part of a larger group of antibodies called antiphospholipid antibodies. An abnormally high level of these antibodies is found in about 15% of women who experience recurrent miscarriage, and is called antiphospholipid syndrome (APS) or Hughes syndrome. (You may also hear it called "sticky blood syndrome".)

## How can these antibodies cause a problem?

It may be that they affect the blood supply in the placenta or that they cause abnormal implantation of the placenta in the wall of the uterus (womb). More research is needed in order to identify exactly how these antibodies cause pregnancy problems.

## Testing

Investigations involve taking a blood sample from the female partner to identify if the antibodies are present. In order to establish a clear diagnosis of antiphospholipid syndrome it is essential to have two positive tests, taken at least six weeks apart.

## Treatment

Treatment is usually with low dose aspirin (75mg daily), starting before conception or early in pregnancy. Your doctor may also recommend injections of low molecular weight heparin once you are pregnant and the baby's heartbeat has been seen on scan.

You may find it helpful to read The Miscarriage Association leaflet *Antiphospholipid syndrome and pregnancy loss*.

## Other immunological investigations

The following tests are less routine, but your doctor may suggest that they are appropriate for you.

### ***Antithrombin III, protein S, protein C, activated protein C resistance (APCR), factor V Leiden and prothrombin gene testings***

It is still not clear whether these factors are linked to miscarriage, but some units test for these at the same time as testing for lupus anticoagulant and anticardiolipin antibodies. If you have an abnormal result, you may be advised to take low dose aspirin and/or heparin. More research is still needed to identify whether this treatment is helpful.

# Hormonal blood tests

## Luteinising Hormone (LH)

### What is this?

LH is a hormone produced by the pituitary gland in the brain. It stimulates a follicle in the ovary containing the egg to burst and release the egg, which then travels from the ovary to the uterus.

### How can LH cause a problem?

In the condition called **Polycystic Ovary Syndrome (PCOS)**, there are usually many small cysts in the ovaries. It is not known why some women have this condition. The incidence of polycystic ovaries is higher in women with recurrent miscarriage and approximately half of these women will produce abnormal amounts of luteinising hormone. Women with high LH levels may find it harder to conceive and are more likely to miscarry when they do conceive. They may also experience problems associated with polycystic ovaries, such as irregular periods, greasy skin and increased body weight.

### Testing

LH levels can be measured in the blood, with the test usually taken between days 2 and 5 of the menstrual cycle. Polycystic ovaries can also be diagnosed by an ultrasound scan.

### Treatment

Despite continuing research into high LH levels and PCOS, there is still no clear and tested treatment for either condition. However, some women may be offered treatment as part of a research trial.

## Progesterone and hCG

Treatment with progesterone and hCG has been tried in an attempt to maintain pregnancy by boosting hormone levels. The scientific evidence is mixed, but your doctor may feel that this treatment is appropriate. More research is being carried out in this area.

## Other hormone and endocrine tests

Some tests performed as part of hospital protocols have never been conclusively linked to miscarriage, unless the levels are very abnormal, and in these cases you would have other symptoms. However, they may still be carried out in some clinics. They include:

### Thyroid Function Test

#### What is this?

The thyroid gland is situated in the neck and produces essential hormones.

#### How can the thyroid cause a problem?

It used to be thought that a thyroid hormone imbalance could cause

miscarriage, but there is currently no clear evidence for this unless the condition is very severe. In this case, however, the symptoms of the thyroid problem would almost certainly have already been investigated and the problem diagnosed. More research is being carried out in this area.

### ***Testing***

Your doctor will take a blood test if there is any suspicion that the level of hormone produced by your thyroid gland is abnormal.

### **Blood sugar level**

Diabetes is not in itself a risk factor for miscarriage, but women whose diabetes is poorly controlled have an increased risk of miscarriage. A routine test for diabetes is not usually performed unless there are symptoms of the condition or a strong family history.

## **Investigation of the uterus (womb)**

It is thought that some cases of miscarriage, especially in later pregnancy, may be due to an abnormal or irregularly-shaped uterus. Sometimes the uterus has an extra wall down its centre, (**bicornuate** or **septate uterus**), or it may have only developed one half (**unicornuate uterus**). It is not clear if such problems cause recurrent miscarriage, but you may be offered tests to investigate this.

### ***Testing***

There are a several ways to investigate the shape of the uterus, including:

- Hysterosalpingogram (HSG)
- Ultrasound
- Saline installation sonography (SIS)
- Hysteroscopy
- MRI Scan

Your doctor will advise you which, if any, seems most appropriate.

### ***Treatment***

If an abnormality of the uterus is found, your doctor may consider surgery, though this will depend upon the type of abnormality. If surgery requires treatment by a specialist, you may have to be referred to another unit. You would always be counselled about whether the abnormality is likely to be causing the miscarriages (it may be coincidental), the risks of treatment, and whether surgery will improve the chance of having a healthy pregnancy. It is always important to weigh up the benefits of treatment with the potential risks of any surgical procedure.

# Investigation of the cervix

## Cervical Resistance Test

This test is sometimes performed if the doctor thinks that there is a weakness of the cervix (sometimes called **cervical incompetence**). There has usually been a history of late miscarriage (i.e. after 14 weeks), with a rapid labour and rupture of the membranes. This condition can be very difficult to diagnose and there is some disagreement about the usefulness of this test.

### **Treatment**

If the test indicates cervical weakness, a cervical stitch may be advised. This procedure is usually carried out under general anaesthetic when a woman reaches 13 or 14 weeks of pregnancy.

*The Miscarriage Association publishes a leaflet on the cervical stitch.*

## Tests for infection

### How can infection cause a problem?

In general, infection is not thought to be a cause of recurrent miscarriage, but severe infection at the time of miscarriage may be the cause of the loss.

There is some evidence that a condition called **bacterial vaginosis (BV)** may cause later miscarriage. In BV, the normal vaginal bacteria are replaced by other bacteria and this causes a vaginal discharge with a 'fishy' smell. Bacterial vaginosis has more recently been associated with early pregnancy loss.

### **Testing**

Your doctor may take a vaginal swab to check for infection and a specific swab for BV. S/he may also suggest blood tests which can indicate if there is an infection.

### **Treatment**

BV is treated with antibiotics.

## Research investigations

There are many hospitals and clinics carrying out research to try to identify why miscarriage occurs. Research may focus on male as well as female factors. If you are asked to take part in a research study, you will be given full information before deciding. Any decision must be yours, but if you feel that you wish to help in the study, then it may benefit you or others who have also suffered from miscarriage.

## Summary

There are many different causes of miscarriage and a number of investigations which can be carried out. In some cases, there may be a combination of causes leading to miscarriage, rather than a single underlying one. Investigations for recurrent miscarriage are usually carried out in a specialised clinic or by a specific consultant in the unit and you may wish to ask your doctor if you can be referred to such a clinic or consultant.

Different clinics will give greater priority to certain investigations. Most centres, however, will test your and your partner's chromosomes and will also test for antiphospholipid antibodies. Many now check height and weight, since very low and very high body mass index (BMI) are both associated with increased risk of miscarriage. The rest of the tests outlined in this leaflet may or may not be required, depending on your medical history.

Don't be afraid to ask questions. The medical and nursing staff will be able to tell you more about the tests they plan to perform, when results will be available and what they mean.

The internet is a source of a great deal of information, but it can be difficult to make sense of different opinions and to judge how accurate this information is. You may read about investigations and treatments for recurrent miscarriage which have not been properly studied or assessed. You may want to discuss what you have read with the team at your recurrent miscarriage clinic.

Finally, it is important to remember that for most couples with a history of recurrent miscarriage, investigations do not identify any specific cause or causes. While this can be very frustrating, it is equally important to remember that for most of you reading this leaflet, you are more likely to have a successful pregnancy next time than to miscarry again. This is especially true if you feel well cared for and supported in your next pregnancy. You might want to talk to your GP and specialist team about how to plan for additional care next time.

If you would like to talk to someone else who has been through the experience of recurrent miscarriage, please contact us at The Miscarriage Association and we will put you in touch with one of our support volunteers.

We hope that this information has been helpful to you.

*Our sincere thanks to Dr Marjory MacLean, Consultant Obstetrician, Ayrshire Maternity Unit, for her help in writing this leaflet; and to GlaxoSmithKline for supporting its production through their Making a Difference programme.*